



**Benjamin B. Bedford, M.D.**  
**159 East 74th Street**  
**New York, NY 10021**  
**Tel (212) 737-3301**

## **Instruction Sheet following Arthroscopic Rotator Cuff Repair Surgery**

### **Sling:**

Wear your sling at **all** times after your surgery (this includes sleeping). There are no exceptions to this rule – it is very important! You will need to wear the sling at all times for about 4 weeks from the date of surgery; this is not related to your level of pain but related to the healing process of the repair. Additionally, you should not carry anything heavier than a pencil in your hand.

### **Dressing:**

Remove all cotton and gauze 48 hours after your surgery. You do not need to put a new dressing on your wound; place a Band-Aid on each of your sutures (stitches).

### **Showering:**

You may shower 48 hours after surgery. Please use **CAUTION!!** Be careful not to slip and fall. The effects of anesthesia and/or medication may make you drowsy or light-headed. Do not submerge your sutures (stitches) in a bathtub, hot tub, or pool until the doctor tells you it is O.K. to do so. Once you are done showering pat the wound dry and apply fresh Band-Aids. While in the shower you must keep the arm across the front of the body as if it were still in the sling, or allow it to gently hang by your side. Be sure not to lift/use the arm to wash yourself.

### **Sleeping:**

You will most likely have difficulty sleeping in the first few weeks after surgery. Most people find it more comfortable to sleep in a reclining position. You can either sleep in a recliner chair or create this position with pillows. You only need to do this if you find it more comfortable.

### **Ice:**

You can ice the shoulder to reduce swelling and discomfort. Do not ice the shoulder more than 20 minutes at a time. Let the shoulder warm up before reapplication of the ice for about 20 minutes. If you have a cryocuff, you may keep this on continuously.

### **Follow-up visit:**



You need to see the doctor about 10-14 days following surgery for your first post-op visit. At that time your sutures (stitches) will be removed, and you will be given a prescription to begin physical therapy.

**Common Concerns:**

Bruising and/or swelling of the shoulder, arm, or hand are common after surgery. This can sometimes extend even into the forearm; this is normal. To relieve this discomfort it is best to ice the shoulder.

**Please call if:**

1. Any oozing or redness of the wound, fevers ( $>101.3^{\circ}\text{F}$ ), or chills.
2. Any difficulty breathing or heaviness in the chest.

**REMEMBER - these are only guidelines for what to expect following arthroscopic rotator cuff repair surgery. If you have any questions or concerns, please do not hesitate to call the office.**